

CROWN AND BRIDGE PROSTHETICS CONSENT FORM

I understand that treatment of dental conditions requiring crowns and/or fixed bridgework includes certain risks and possible unsuccessful results, with even the possibility of failure. I agree to assume those risks, possible unsuccessful results and/or failure associated with, but not limited to the following: (Even though care and diligence is exercised in the treatment of conditions requiring crowns, bridgework and fabrication of same, there are no promises or guarantees of anticipated results or longevity of treatment.)

Reduction of tooth structure: In order to replace decayed or otherwise traumatized teeth it is necessary to modify the existing tooth or teeth so that crowns (caps) and/or bridges may be placed upon them. Tooth preparation will be done as conservatively as practical. In preparation of teeth, anesthetics are usually needed. At times there may be swelling, jaw muscle tenderness or even a resultant numbness of the tongue, lips, teeth, jaws and/or facial tissues which is usually temporary, or rarely permanent.

Sensitivity of teeth: Often, after the preparation of teeth for the reception of either crowns or bridges, the teeth may exhibit sensitivity. It may be mild to severe. The sensitivity may last only for a short period of time or may last for much longer periods. If it is persistent, notify us immediately, as this sensitivity may be from some other source.

Crowned or bridge abutment teeth may require root canal treatment: Teeth after being crowned may develop a condition known as pulpitis or pulpal degeneration. The tooth or teeth may have been traumatized from an accident, deep decay, extensive preparation, or other causes. It is often necessary to do root canal treatments in these teeth. If teeth remain too sensitive for long periods of time following crowning, root canal treatment may be necessary. Infrequently, the tooth/teeth may abscess or otherwise not heal which may require root canal treatment, root surgery, or possibly extraction.

Breakage: Crowns and bridges may possibly chip or break. Many factors could contribute to this situation such as chewing excessively hard materials, changes in biting forces, traumatic blows to the mouth, etc. Unobservable cracks may develop in crowns from the above-mentioned factors. Crowns and bridges may inadvertently break secondary to unobservable cracks. Breakage or chipping seldom occurs due to defective materials or construction unless it occurs soon after placement.

Uncomfortable or strange feeling: This may occur because of the differences between natural teeth and the artificial replacements. Most patients usually become accustomed to this feeling in time. In limited situations, muscle soreness or tenderness of the jaw joints (TMJ) may persist for indeterminable periods of time following placement of prosthesis.



Esthetics or appearance: I understand that sometime it is not possible to match the color of natural teeth exactly with artificial teeth. I further understand that I may be wearing temporary crowns, which may come off easily and that I must be careful to ensure that they are kept on

until the permanent crowns are delivered. Patients will be given the opportunity to observe the appearance of crowns or bridges in place prior to final cementation. I realize that final opportunity to make changes in my new crown, bridge, or prosthetic (including shape, fit, size, and color) will be before cementation.

Longevity of crowns and bridges: There are many variables that determine "how long" crowns or bridges can be expected to last. Among these are some of the factors mentioned in preceding paragraphs. Additionally, general health, good oral hygiene, regular dental checkups, diet, etc., can affect longevity. Because of this, no guarantees can be made or assumed to be made.

It is the patient's responsibility to seek attention from the dentist should any undue or unexpected problems occur. The patient must diligently follow any and all instructions, including the scheduling and attending of all appointments. Failure to keep the cementation appointment can result in ultimate failure of the crown/bridge to fit properly and an additional fee may be assessed.

Informed consent: I have been given the opportunity to ask any questions regarding the nature and purpose of crown and/or bridge treatment and have received answers to my satisfaction. I voluntarily assume any and all possible risks including those listed above and including risk of substantial harm, if any, which may be associated with any phase of this treatment in hopes of obtaining the desired results, which may or may not be achieved. No guarantees or promises have been made to me concerning the results. The fee(s) for service have been explained to me and are satisfactory. By signing this document, I am freely giving my consent to allow Dlight Dental to render any treatment necessary and/or advisable to my dental conditions including the prescribing and administering any medications and/or anesthetics deemed necessary to my treatment.

Patient Signature/Legally authorized representative

Print Name

Date

