

Surgery Follow Up

After any surgery, minor discomfort, bleeding and swelling is normal. The following instructions have been reviewed with you by your doctor and are for your reference.

DISCOMFORT: After any surgical procedure a certain amount of discomfort is expected. By taking the prescribed medications as directed, your discomfort should be reduced. Under no circumstances should you take Aspirin for pain, since it encourages bleeding. Teeth that were treated may become more mobile or sensitive to hot, cold and brushing. These problems are usually temporary but please inform your doctor if it becomes problematic.

BLEEDING: Expect some pink fluid in your mouth. This is simply saliva mixing with the blood clot, and it is of no significance. You may want to place a towel over your pillow the night after surgery to protect your bed linen. Slight oozing from the area of surgery is normal on the first day of surgery. Because blood mixes with saliva, it may appear to be a greater volume than it really is. To manage post---operative bleeding you should:

1. Avoid rinsing, spitting or using a straw for 24 hours
2. Avoid alcohol and smoke for at least 72 hours
3. Avoid vigorous exercise for 72 hours
4. Avoid eating hard or sharp foods that may further traumatize the wound

If bleeding does occur, apply pressure to area for twenty minutes with a piece of DAMP gauze. If bleeding continues apply pressure for an additional twenty minutes with a DAMP tea bag (regular tea, not herbal). If bleeding is severe and does not stop, please contact your doctor.

SWELLING: Swelling usually occurs 1-2 days after surgery. To minimize swelling apply ice for 20 minutes on and 20 minutes off for the first 24 hours. Sleep with an extra pillow under your head for the first night and take the anti-inflammatory medications recommended by your doctor. Some bruising may occur and should not alarm you. You may also experience stiffness of the jaw which can be relieved by gentle opening and stretching on the 2nd and 3rd day after surgery.

DIET: Following surgery you may experience mild tooth discomfort, and as such, maintaining a softer diet is strongly recommended. Foods such as overcooked pasta, mashed potatoes, eggs and soups are a well-balanced option. Avoid foods:

1. With small pieces, such as nuts, fruits with seeds, chips etc. can be trapped in the surgical area
2. That is too hot, or too spicy for 24 hours
3. With alcohol as it may interact with medications you are taking

It is important to eat a full but modified diet as your body needs proteins, vitamins, minerals and calories to heal and fight infection.

ORAL HYGIENE: It is very important to keep your mouth as clean as possible to reduce the chance of infection. 24hours after the surgery: no rinsing, hence no brushing of teeth. Resume your routine oral hygiene, including brushing on unaffected areas and rinsing 24 hours after the surgery. Avoid flossing the surgical area 2 month or until asked by doctor to resume it.

REST: After surgery, you should rest as much as possible for 1-2 days to allow your body to heal. Your post-operative comfort is very important to us. If you have any concerns, please do not hesitate to contact our office during business hours or Doctor on 617-903-7838 after business hours.

